

*Variability of Low-Level Temperature Inversions in California:  
Applying GCM Results to California Air Basins*


Sam Iacobellis

*Collaborators:*

Dan Cayan, Masao Kanamitsu, Joel Norris

Climate Research Division  
Scripps Institution of Oceanography  
University of California, San Diego

Project Funded by California Air Resources Board




Mean temperature profile at San Diego (KSAN)  
as a function of time of day.

## Outline


1. Numerical models
2. Low-level temperature inversions
3. Relationship of inversion strength to large-scale and regional-scale circulation

# General Circulation Models (GCMs)

- System of equations describing atmosphere and ocean
- Relatively coarse grid  $\sim 2.5^\circ \times 2.5^\circ$
- Models used to make future global climate predictions (IPCC report)



**Elevation**




meters




0 1000 2000 3000 4000

**Elevation**




**NCEP-R2 2.5 Degree Resolution**



meters




meters




## Downscaling

- Provide increased horizontal and vertical resolution
- Regional instead of global
- Use coarse global model results as boundary conditions




Regional Dynamical Model at Scripps: CaRD10 - California Reanalysis Downscaling at 10 km  
Statistical models also being developed


**Elevation**




meters




**NCEP-R2 2.5 Degree Resolution**




meters



**CaRD10 10 km Resolution**




meters





# Temperature Inversions



Mean temperature profile at San Diego  
as a function of time of day.

# Temperature Inversions

Possible Measures:


$$DTINV = T_{\text{top}} - T_{\text{base}}$$

$$DT850 = T_{850} - T_{2m}$$

T850 = Temperature at 850 mb


DTDZ = lapse rate within inversion

PBASE = Inversion base pressure




Mean temperature profile at San Diego  
as a function of time of day.

## Inversions vary seasonally, but are a dominant feature in California air basins





# Temperature inversions and pollution



# Temperature inversions and pollution

**Correlation of Daily Means  
Inversion Measure vs Pollutant  
Month = June**



## Relationship of inversion strength to large-scale and regional-scale circulation


Model data from:

- NCEP Reanalysis 2 ( $2.5^\circ \times 2.5^\circ$ )
  - similar resolution to most climate models
  - hindcast
  - incorporates available observations
  - represents best estimate of atmospheric state 1979-present
- California Reanalysis Downscaling at 10km (CaRD10)
  - dynamical downscaling


## Composite Daily Atmospheric Patterns During Strong/Weak Inversion Events

- examine weather balloon data at Oakland (Jun-Aug 1979-2001)
- find the 30 events with largest/smallest inversion magnitudes
- examine mean large-scale circulation for these 30 events
- consider anomalies (departure from long-term average)

## 500mb Height and Wind Anomalies



Strong Inversions  
at Oakland



Weak Inversions  
at Oakland


- Strong inversions associated with above normal 500mb heights (large-scale high pressure systems)
- Weak inversions associated with below normal 500mb heights (large-scale low pressure systems)

==> Inversions in California associated with large-scale circulation




**DOWNSCALED COMPOSITE MEANS JUN-AUG  
SURFACE WIND AND INVERSION MAGNITUDE  
ACTUAL VALUES (NOT ANOMALIES)**


**STRONG INVERSIONS AT OAKLAND**



**WEAK INVERSIONS AT OAKLAND**



## Large-Scale 500mb Height Difference



Define  $DH500 = H_{500,reg1} - H_{500,reg2}$  using historical analysis data



How does this large-scale variable relate to local inversion measures in throughout California? On daily timescales? Monthly timescales?

# HOW DO LOCAL INVERSION MAGNITUDES COMPARE TO LARGE-SCALE FEATURES?



## CORRELATION OF DAILY MEANS

Downscaled Inversion Magnitude vs. Large-Scale 500mb Height Difference

MAR-APR-MAY 1979-2005



JUN-JUL-AUG 1979-2005



## California Inversion Index GFDL A2

500hPa height diff, Elko minus Churchill




Figure 9. Frequency (5-year running total) of positive  $^2$  h500 anomalies exceeding 1.0 standard deviations from the SRES A2 runs of the GFDL CM2.1 model. Here  $^2$ h 500 is defined as the difference in 500 mb height between 42°N, 115°W (Elko) and 60°N, 95°W (Churchill). The anomalies are referenced to the 1961-1990 climatology.

## Summary

- Low-level temperature inversions are a consistent feature in California air basins as evidenced at San Diego and Oakland (additional soundings at inland sites currently being examined).
- Inversion measures using temperature at top or across inversion show higher correlation to pollution.
- Inversion intensity is strongly linked to large-scale circulation features (e.g., CA Central Valley during Spring). This large-scale structure should allow us to downscale GCM simulations to California air basins.
- Climate change simulations provide large-scale structure that may allow us to project trends of inversion magnitudes and frequencies.

## Future Work

- Continue to collect and analyze atmospheric soundings
- Produce more quantitative products relating large-scale circulation to inversion characteristics for California air basins
- Examine how inversion frequency/magnitude related to warm/cold periods in historical record
- Investigate role of decadal climate modes and ENSO on circulation patterns and inversion characteristics
- Use GCM climate simulations and downscaling to examine potential changes in low-level inversions.